

Shigella

Your Health. Our Work.

www.GCHD.us

What is shigella?

• A bacterium that causes a diarrheal disease. *Shigella* causes about 500,000 cases of diarrhea in the United States annually. Symptoms normally occur within 1-2 days after being infected with the bacteria and can last 5-7 days.

What are symptoms that could occur?

Diarrhea

Fever

- Abdominal pain
- Tenesmus (painful sensation of needing to pass stools even when bowels are empty)

How is it spread?

- Contaminated hands or surfaces touch your food or mouth (toys, bathroom fixtures, changing tables, diapers)
- Eating food contaminated with shigella that was prepared by food handlers who were infected
- Swallowing recreational or drinking water that was contaminated by infected fecal matter
- Exposure to feces through sexual contact

How is it diagnosed and treated?

- Primary care provider evaluation and laboratory evaluation
- The laboratory can also do special tests to determine if antibiotics are best to treat the infection

Are there complications?

- Diarrhea and vomiting may cause the body to lose a lot of fluids, causing dehydration
- Rarely, post-infectious arthritis, blood stream infections, seizures, and hemolytic-uremic syndrome (HUS) may occur

How can shigella be prevented?

- Carefully wash your and your child's hands with soap and water often
- If you or your child is sick, stay home from work or school
- If you care for a child in diapers who has *shigella*, promptly discard the soiled diapers in a lidded/lined garbage can, and wash your hands and the child's hands carefully with soap and water immediately after changing the diaper
- Avoid swallowing water from ponds, lakes or untreated swimming pools
- Avoid sexual activity with those who have diarrhea or who recently recovered from bacteria

Should the sick individual be isolated?

• Yes. Infected person should be kept away from other people until person feels well again.

When can the individual return to the general population?

• When the person feels well again and diarrhea has stopped for at least 48 hours. Food handlers and healthcare workers should wait until diarrhea has stopped for at least 72 hours.

Should facilities do special cleaning

• Special cleaning is not necessary. Routine cleaning with EPA approved products is recommended

If I have any other questions, whom can I contact?

Call the Genesee County Health Department at 257-1017

This fact sheet is for information only and is not meant to be used for self-diagnosis or as a substitute for consultation with a health care provider. For more information, call your health care provider. 01/2016